

Structural Investigation of Large Zincferrocenedicarboxylate Clusters

A.Tripathi, and J.B. Parise (SUNY, Stony Brook), O.M.Yaghi and H. Li (U. Michigan)

Abstract No. Trip0504

Beamline(s): X3A1

Introduction: A recent new approach in the preparation of robust framework involves the employment of bis- and tris bidentate organic linkers such as 1,4-benzenedicarboxylate [1]. In this series, dark red crystals of zincferrocenedicarboxylate have been synthesized and characterized by single crystal synchrotron X-ray diffraction. $\text{Zn}_3(\text{Fec})_3(\text{DMF})_2(\text{H}_2\text{O})_3$ (where FecH_2 is 1,1'-Ferrocenedicarboxylic acid and DMF is Dimethylformamide) is monoclinic, space group $P2_1/n$ with $a = 14.990(1)$, $b = 22.606(2)$, $c = 22.994(1)$ Å, $\beta = 98.010(1)^\circ$. The structure consisting of 118 atoms has two types of Zn coordination, four zinc sites have tetrahedral and two have octahedral coordination. Each cluster consists of Zn octahedra sharing a corner with two other Zn tetrahedra. Two such units are capped by six-ferrocene dicarboxylate in a bidentate fashion. The clusters are held in position by hydrogen bonds with water molecules and DMF.

Acknowledgments: The authors thank the NSF for financial support (Grant DMR 97-13375). Research carried out in part at the NSLS at BNL is supported by the U.S. Department of Energy, Division of Materials Sciences and Division of Chemical Sciences, Office of Basic Energy Sciences (Grant DE-FG02-86ER45231 for the SUNY X3A beamline).

References: [1] M. Eddaoudi, H.Li, and O.M. Yaghi. (2000) J. Am. Chem. Soc., **122**, 1391




Figure 1. Polyhedral representation of zincferrocenedicarboxylate clusters. 1,1' ferrocenedicarboxylate units (purple and green) cap yellow tetrahedra and octahedra of zinc. The ferrocene units are shown here with 2 carbons attached to an iron for clarity. Blue spheres represent water and DMF, bonded to these clusters by H-bonds. The red spheres represent oxygen atoms.